

MATHICSE Technical Report

Nr. 05.2013

February 2013 (NEW 10.2013)



A weighted empirical interpolation method: a priori convergence analysis and applications

P. Chen, A. Quarteroni, G. Rozza

A WEIGHTED EMPIRICAL INTERPOLATION METHOD: A PRIORI CONVERGENCE ANALYSIS AND APPLICATIONS

PENG CHEN¹ · ALFIO QUARTERONI^{1 2} · GIANLUIGI ROZZA³

Abstract: We extend the classical empirical interpolation method [1] to a weighted empirical interpolation method in order to approximate nonlinear parametric functions with weighted parameters, e.g. random variables obeying various probability distributions. A priori convergence analysis is provided for the proposed method and the error bound by Kolmogorov N-width is improved from the recent work [13]. We apply our method to geometric Brownian motion, exponential Karhunen-Loève expansion and reduced basis approximation of non-affine stochastic elliptic equations. We demonstrate its improved accuracy and efficiency over the empirical interpolation method, as well as sparse grid stochastic collocation method.

Keywords: empirical interpolation method, a priori convergence analysis, greedy algorithm, Kolmogorov N-width, geometric Brownian motion, Karhunen-Loève expansion, reduced basis method

Mathematics Subject Classification. 65C20, 65D05, 97N50

1 Introduction

The empirical interpolation method [1] was originally developed to approximate the non-affine terms of a partial differential equation in order to effectively decompose the reduced basis method into offline construction and online evaluation procedure. Since its development, many applications and extensions of this method have been considered [10, 13, 22, 20, 12, 4, 18]. In particular, we mention its application and analysis in the context of reduced basis approximation for nonlinear elliptic and parabolic equations [10] and, more recently, its extension to a general, multipurpose interpolation procedure [13], in which a priori error estimate compared to Kolmogorov N-width was obtained.

The basic idea behind empirical interpolation for parametric function $g(x, \mu)$ is to choose the parameter samples μ^1, μ^2, \dots and the interpolation nodes x^1, x^2, \dots recursively in a greedy approach according to the criteria that μ^m and x^m selected at each step $m = 1, 2, \dots$ are the most representative ones in L^∞ norm or the ones where the function is worst approximated by the interpolation formula constructed from the previous steps [1]. This is essentially different from the conventional interpolation construction which requires the interpolation nodes to be chosen a priori according to a specific rule, e.g. roots of orthogonal polynomials [19]. The so called “magic points” [13] $(\mu^m, x^m), m = 1, 2, \dots$ obtained by the goal-oriented or function-specified empirical interpolation procedure are supposed to identify an interpolation formula by capturing some specific features (e.g. regularity, extreme values) of

¹Modelling and Scientific Computing, CMCS, Mathematics Institute of Computational Science and Engineering, MATHICSE, Ecole Polytechnique Fédérale de Lausanne, EPFL, Station 8, CH-1015 Lausanne, Switzerland. Peng Chen (peng.chen@epfl.ch, cpempire@gmail.com), Alfio Quarteroni (alfio.quarteroni@epfl.ch)

²Modellistica e Calcolo Scientifico, MOX, Dipartimento di Matematica F. Brioschi, Politecnico di Milano, P.za Leonardo da Vinci 32, I-20133, Milano, Italy

³SISSA MathLab, International School for Advanced Studies, via Bonomea 265, 34136 Trieste, Italy. (gianluigi.rozza@sissa.it)

the given function, thus providing higher interpolation accuracy. Another superiority of the empirical interpolation construction is attributed to the affine expansion of the function given in whatever form, leading to the separation of the variable x and the parameter μ in the following expression [10]

$$g(x, \mu) \approx \mathcal{I}_M[g] = \sum_{j=1}^M \Theta_j(\mu) q_j(x), \quad (1.1)$$

which can be efficiently used in conducting mathematical manipulation, e.g. numerical integration, reduced basis approximation [10]. By convention, one supposes that the parameter μ , if viewed as a random variable, is uniformly distributed in a bounded space Γ .

However, in many applications, e.g. stochastic problems with parametrized random variables obeying normal distribution, the request of the boundedness of the parameter space Γ and that of the uniform distribution of the parameter μ is hard to fulfill. In this situation, the approximation to some quantity of interest (e.g. weighted integral or statistics of the function) based on the parameter samples and interpolation nodes selected by the empirical interpolation procedure would not lead to results that are as accurate or efficient as those expected when taking distinct weights of the parameter at different values into account. In this work we propose a weighted empirical interpolation method (wEIM) by considering a weighted optimization problem and analyzing its convergence property by improving the a priori error estimate obtained in [13]. To demonstrate numerically its effectiveness and efficiency, we apply the wEIM to approximating nonlinear parametric functions, geometric Brownian motion in one dimension, exponential Karhunen-Loève expansion in multi-dimension as well as reduced basis approximation to non-affine stochastic elliptic problems, and compare it with the conventional empirical interpolation method (EIM) and sparse grid stochastic collocation method. It is worth mentioning that constructing a goal-oriented numerical method is a quite common procedure in adaptive finite element methods [2, 9] and has also been applied to construct adaptive reduced basis method [5].

The work is organized as follows: we present the weighted empirical interpolation method in section 2. A priori convergence analysis is carried out in section 3, followed by section 4 where different applications of this method are addressed. Some concluding remarks are drawn in section 5.

2 Weighted empirical interpolation method (wEIM)

For notational convenience, we introduce the spaces $L^\infty(D)$ defined in a bounded physical domain $D \subset \mathbb{R}^d, d \in \mathbb{N}_+$ and $C_w^0(\Gamma)$ defined in a parameter space (not necessarily bounded) $\Gamma \subset \mathbb{R}^K, K \in \mathbb{N}_+$, which are equipped with the following norms: $\|g\|_{L^\infty(D)} = \text{ess sup}_{x \in D} |g(x)|$ and $\|g\|_{C_w^0(\Gamma)} = \max_{\mu \in \Gamma} w(\mu) |g(\mu)|$ with a positive weight function $w : \Gamma \rightarrow \mathbb{R}_+$. We also define the Bochner space $L^\infty(D; C_w^0(\Gamma))$ for a parameter dependent function equipped with the norm $\|g\|_{L^\infty(D; C_w^0(\Gamma))} = \text{ess sup}_{x \in D} (\max_{\mu \in \Gamma} w(\mu) |g(x, \mu)|) \equiv \max_{\mu \in \Gamma} w(\mu) (\text{ess sup}_{x \in D} |g(x, \mu)|)$. We note that $L^\infty(D)$, as used in [1, 10, 13], is usually replaced with $C^0(D)$ for conventional interpolation [19].

At the discrete level, the physical domain D is replaced by a series of vertices $x \in V_x$ with finite cardinality $n_x = |V_x| < \infty$, for instance finite element nodes, and the parameter space Γ is represented by a sample set $\mu \in \Xi_\mu$ of finite cardinality $n_\mu = |\Xi_\mu| < \infty$. We present the weighted empirical interpolation method in the following generic Algorithm 1. We emphasize that the initial sample μ^1 is chosen such that the weighted function is maximized in $L^\infty(V_x; C_w^0(\Xi_\mu))$ norm:

$$\mu^1 = \arg \max_{\mu \in \Xi_\mu} \left[w(\mu) \left(\text{ess sup}_{x \in V_x} |g(x, \mu)| \right) \right]. \quad (2.1)$$

In the course of the construction procedure, the quasi-optimal samples $\mu^{M+1}, M \geq 1$ can be chosen by greedy algorithm to minimize the weighted optimal approximation error (2.2) in the subspace W_M spanned by the ‘‘snapshots’’ $W_M := \text{span}\{g(\cdot, \mu^i), 1 \leq i \leq M\}$: find $\mu^{M+1} \in \Xi_\mu$ such that

$$\mu^{M+1} = \arg \max_{\mu \in \Xi_\mu} \left[w(\mu) \left(\inf_{h \in W_M} \|g(\mu) - h\|_{L^\infty(V_x)} \right) \right]. \quad (2.2)$$

However, the weighted L^∞ optimization problem (2.2) is expensive to solve by linear programming if $|V_x|$ and $|\Xi_\mu|$ are large. In practice, it can be efficiently replaced by a weighted L^2 optimization problem [10] or by a surrogate weighted L^∞ optimization problem (2.5) [13].

Algorithm 1 A weighted empirical interpolation method

1: **procedure** INITIALIZATION:

2: Given finite vertex set $V_x \subset D$, sample set $\Xi_\mu \subset \Gamma$, weight w and function $g \in L^\infty(V_x; C_w^0(\Xi_\mu))$;

3: find $\mu^1 \in \Xi_\mu$ such that $\mu^1 = \arg \max_{\mu \in \Xi_\mu} w(\mu)(\text{ess sup}_{x \in V_x} |g(x, \mu)|)$; set $W_1 = \text{span}\{g(x, \mu^1)\}$;

4: find $x^1 \in V_x$ such that $x^1 = \arg \text{ess sup}_{x \in V_x} |g(x, \mu^1)|$;

5: define $r_1 = wg$, $q_1(x) = r_1(x, \mu^1)/r_1(x^1, \mu^1)$, $B_{11}^1 = 1$, set $M = 1$, specify tolerance ε_{tol} ;

6: **end procedure**

7: **procedure** CONSTRUCTION:

8: **while** $M < M_{max}$ & $r_M(x^M, \mu^M) > \varepsilon_{tol}$ **do**

9: find $\Theta^M(\mu^M) = (\Theta_1^M(\mu^M), \dots, \Theta_M^M(\mu^M))^T$ by solving

$$\sum_{j=1}^M \Theta_j^M(\mu^M) q_j(x^i) = g(x^i, \mu^M) \quad 1 \leq i \leq M; \quad (2.3)$$

10: define $r_{M+1} : D \times \Gamma \rightarrow \mathbb{R}$ as

$$r_{M+1}(x, \mu) = g(x, \mu) - \sum_{j=1}^M \Theta_j^M(\mu) q_j(x); \quad (2.4)$$

11: find $\mu^{M+1} \in \Xi_\mu$ such that

$$\mu^{M+1} = \arg \max_{\mu \in \Xi_\mu} \left[w(\mu) \left(\text{ess sup}_{x \in V_x} |r_{M+1}(x, \mu)| \right) \right], \quad (2.5)$$

12: find $x^{M+1} \in V_x$ such that

$$x^{M+1} = \arg \text{ess sup}_{x \in V_x} |r_{M+1}(x, \mu^{M+1})|; \quad (2.6)$$

13: define $q_{M+1} : D \rightarrow \mathbb{R}$ as

$$q_{M+1}(x) = \frac{r_{M+1}(x, \mu^{M+1})}{r_{M+1}(x^{M+1}, \mu^{M+1})}; \quad (2.7)$$

14: update matrix $B^{M+1} \in \mathbb{R}^{(M+1) \times (M+1)}$ as

$$B_{ij}^{M+1} = q_j(x^i) \quad 1 \leq i, j \leq M+1; \quad (2.8)$$

15: **end while**

16: **end procedure**

17: **procedure** EVALUATION:

18: For $\forall \mu \in \Xi_\mu$, construct approximation (1.1) by solving (2.3), then evaluate (1.1) at $\forall x \in V_x$.

19: **end procedure**

We state several properties of the wEIM in the following lemmas, whose proof is straightforward by noting the fact that the weight function $w : \Gamma \rightarrow \mathbb{R}_+$ is positive and therefore omitted here, see for instance [1, 10, 13] for details.

Lemma 2.1 *For any $M < M_{max}$, the subspace $Q_M = \text{span}\{q_m, 1 \leq m \leq M\}$ is of dimension M . Moreover, the matrix B^M formed in (2.8) is lower triangular with unity diagonal and thus invertible.*

Lemma 2.2 *For any function $h \in Q_M$, the empirical interpolation formula given by (2.3) is exact,*

i.e. $r_{M+1}(x, \mu) = 0, \forall x \in V_x, \mu \in \Xi_\mu$. In general, for any function $g \in L^\infty(D; C_w^0(\Gamma))$, we have

$$\|g - \mathcal{I}_M[g]\|_{L^\infty(D)} \leq (1 + \Lambda_M) \inf_{h \in Q_M} \|g - h\|_{L^\infty(D)} \text{ with } \Lambda_M \leq 2^M - 1. \quad (2.9)$$

3 A priori convergence analysis

The interpolation error obtained in (2.9) with the Lebesgue constant $\Lambda_M \leq 2^M - 1$ (see proof in [10]) by the empirical interpolation procedure is too pessimistic, far from the result for conventional interpolation error based on certain prescribed interpolation nodes (e.g. Chebyshev nodes) with $\Lambda_M \sim \log(M)$ [19]. An explicit a priori convergence rate of the weighted empirical interpolation error is not available for generic functions. In order to measure the accuracy of the approximation by wEIM, in the following theorem we compare it with the Kolmogorov N -width [17], which quantifies the optimal approximation error of a subset \mathcal{F} in a Banach space \mathcal{H} by any possible N dimensional subspace F_N , defined as

$$d_N(\mathcal{F}, \mathcal{H}) := \inf_{F_N} \sup_{g \in \mathcal{F}} \inf_{f \in F_N} \|g - f\|_{\mathcal{H}}. \quad (3.1)$$

When the subset \mathcal{F} becomes the same as \mathcal{H} , we denote $d_N(\mathcal{H}) \equiv d_N(\mathcal{F}, \mathcal{H})$ for simplicity.

Theorem 3.1 *The error of wEIM can be bounded as follows*

$$\|g - \mathcal{I}_M[g]\|_{L^\infty(V_x)} \leq C_w(M+1)2^M d_M(L^\infty(V_x)) \quad (3.2)$$

where the constant C_w depends on the weight function w but is independent of M .

Remark 3.1 *In fact, the result (3.2) is obtained in the subspace $L^\infty(V_x)$ for the constructive wEIM and can be straightforwardly extended to $L^\infty(D)$ when the vertex set V_x tends to D such that the points outside the vertex set V_x can be sufficiently well represented by the points inside. To be rigorous, we take the vertex set V_x such that for almost every $x \in D$, there exists $y \in V_x$ satisfying*

$$|g(x) - g(y)| \leq \|g - \mathcal{I}_M[g]\|_{L^\infty(V_x)}. \quad (3.3)$$

Consequently, we have the error bound

$$\begin{aligned} \|g - \mathcal{I}_M[g]\|_{L^\infty(D)} &\leq \|g - \mathcal{I}_M[g]\|_{L^\infty(D \setminus V_x)} + \|g - \mathcal{I}_M[g]\|_{L^\infty(V_x)} \\ &\leq 2\|g - \mathcal{I}_M[g]\|_{L^\infty(V_x)} \\ &\leq C_w(M+1)2^{M+1}d_M(L^\infty(V_x)) \\ &\leq C_w(M+1)2^{M+1}d_M(L^\infty(D)). \end{aligned} \quad (3.4)$$

The proof of (3.2) adopts a constructive approach inspired from that for the greedy algorithm in reduced basis approximation [3, 21]. Some preliminary results are needed, see the next two lemmas.

For simplicity, we use the shorthand notation $r_m(x) = r_m(x, \mu^m), 1 \leq m \leq M+1$ obtained in Algorithm 1 and define the functions $t_j(x^i) = r_i(x^j), 1 \leq i, j \leq M+1$ and $t_j(x^i) = 0, 1 \leq j \leq M+1, i > M+1$.

Lemma 3.2 *The matrix T^{M+1} defined by $T_{ij}^{M+1} = t_j(x^i), 1 \leq i, j \leq M+1$ is upper triangular matrix with dominating diagonal elements, i.e. $t_j(x^i) = 0, i > j$ and $|t_j(x^i)| \leq |t_j(x^j)|, i \leq j$.*

Proof From the result of Lemma (2.1), we know that the matrix B^{M+1} is lower triangular with unity diagonal. By the definition of $q_i, 1 \leq i \leq M+1$ in (2.7) and the definition of $t_j, 1 \leq j \leq M+1$, we have $t_j(x^i) = q_i(x^j)r_i(x^i)$, so that $t_j(x^i) = q_i(x^j) = 0, i > j$ and $|t_j(x^i)| \leq |t_j(x^j)| = |r_j(x^j)|, i \leq j$ due to (2.6). \square

Lemma 3.3 For any $1 \leq m \leq M + 1$, there exists a unique $b = (b_1, \dots, b_m)^T \in \mathbb{R}^m$ such that

$$r_m(x)e_m(x) = \sum_{j=1}^m b_j t_j(x) \quad \forall x \in V_x, \quad (3.5)$$

where $e_m, 1 \leq m \leq M + 1$ are unit vectors, i.e. $e_m(x^m) = 1$ and $e_m(x^n) = 0$ if $n \neq m$. In addition, we have $b_m = 1$ and the bound $|b_i| \leq 2^{m-i-1}, 1 \leq i < m$ so that $|b_1| + \dots + |b_m| \leq 2^{m-1}$.

Proof For any $x = x^i, i > M + 1$, we have $e_m(x) = 0$ and $t_j(x) = 0$, so that both sides of the equation vanish and we only need to verify the statement for $x = x^i, 1 \leq i \leq M + 1$, in which case the system (3.5) becomes

$$s = Tb \quad \text{with } s = (0, \dots, 0, r_m(x^m))^T. \quad (3.6)$$

Thanks to Lemma 3.2, we have that T is invertible and thus there exists a unique solution b . Moreover, the last line of the system (3.6) $r_m(x^m) = t_m(x^m)b_m$ leads to the solution $b_m = 1$ since $r_m(x^m) = t_m(x^m)$. For any other line $i, 1 \leq i < m$, we have by the fact that T is an upper triangular matrix

$$0 = \sum_{j=i}^m b_j t_j(x^i). \quad (3.7)$$

By recalling that $|t_j(x^i)| \leq |t_i(x^i)|, j > i$, this yields the following bound for $b_i, 1 \leq i < m$

$$|b_i| = \left| - \sum_{j=i+1}^m b_j \frac{t_j(x^i)}{t_i(x^i)} \right| \leq \sum_{j=i+1}^m |b_j|, \quad (3.8)$$

so that $|b_i| \leq 2^{m-i-1}, 1 \leq i < m$ and $|b_1| + \dots + |b_m| \leq 2^{m-1}$ being $b_m = 1$ and using a recursive argument. \square

We are now ready to prove Theorem 3.1 using the representation of the residual in Lemma 3.3.

Proof of Theorem 3.1. Suppose there exists a subspace $H_M \subset L^\infty(V_x)$ of dimension M achieving the Kolmogorov M -width as defined in (3.1), then we have a series of elements $h_j \in H_M, 1 \leq j \leq M + 1$ such that

$$\|t_j - h_j\|_{L^\infty(V_x)} \leq d_M(L^\infty(V_x)), 1 \leq j \leq M + 1. \quad (3.9)$$

We define the functions

$$s_m(x) = \sum_{j=1}^m b_j h_j(x), 1 \leq m \leq M + 1. \quad (3.10)$$

Since all the elements $h_j, 1 \leq j \leq M + 1$ belong to the M dimensional subspace H_M and s_m is a linear combination of these elements for any $m = 1, \dots, M$, there exists a vector $\alpha = (\alpha_1, \dots, \alpha_{M+1})^T$ with $|\alpha_1| + \dots + |\alpha_{M+1}| = 1$ such that

$$\sum_{m=1}^{M+1} \alpha_m s_m = 0. \quad (3.11)$$

Thanks to the result in Lemma 3.3 together with bound (3.9) and representation (3.10) and (3.11), we obtain the following bound for every $x \in V_x$

$$\begin{aligned}
\left| \sum_{m=1}^{M+1} \alpha_m r_m(x) e_m(x) \right| &= \left| \sum_{m=1}^{M+1} \alpha_m (r_m(x) e_m(x) - s_m(x)) \right| \\
&\leq \left(\sum_{m=1}^{M+1} |\alpha_m| \right) \max_{m=1, \dots, M+1} \|r_m e_m - s_m\|_{L^\infty(V_x)} \\
&\leq \max_{m=1, \dots, M+1} \left(\sum_{j=1}^m |b_j| \right) \max_{j=1, \dots, m} \|t_j - h_j\|_{L^\infty(V_x)} \\
&\leq 2^M d_M(L^\infty(V_x)).
\end{aligned} \tag{3.12}$$

Since $|\alpha_1| + \dots + |\alpha_{M+1}| = 1$, there must exist α_m such that $|\alpha_m| \geq 1/(M+1)$. Setting $x = x^m$ in (3.12), we have $|\alpha_m r_m(x^m)| \leq 2^M d_M(L^\infty(V_x))$ and thus

$$|r_m(x^m)| \leq (M+1)2^M d_M(L^\infty(V_x)). \tag{3.13}$$

By the construction of weighted empirical interpolation approximation in Algorithm 1, we have

$$\operatorname{ess\,sup}_{x \in V_x} |r_{M+1}(x)| \leq |r_{M+1}(x^{M+1})| \leq |r_M(x^M)| \leq \dots \leq |r_m(x^m)|. \tag{3.14}$$

A combination of (3.13) and (3.14) leads to the following error bound

$$\|g - \mathcal{I}_M[g]\|_{L^\infty(V_x)} \leq \operatorname{ess\,sup}_{x \in V_x} |r_{M+1}(x)| \leq (M+1)2^M d_M(L^\infty(V_x)). \tag{3.15}$$

□

Corollary 3.4 *Under the assumption $d_M(L^\infty(V_x)) \leq ce^{-rM}$ being $r > \log(2)$, we have the following a priori error estimate of the wEIM: for $\forall g \in L^\infty(V_x; C_w^0(\Xi_\mu))$*

$$\|g - \mathcal{I}_M[g]\|_{L^\infty(V_x)} \leq c(M+1)e^{-(r-\log(2))M}. \tag{3.16}$$

Remark 3.2 *The result (3.16) is an improvement of that recently obtained in [13], in which r is required to satisfy $r > 2\log(2)$ and the exponential convergence rate becomes $r - 2\log(2)$. In fact, when the function g is analytic with respect to the parameter $\mu \in \mathbb{R}$, the Kolmogorov width is bounded by the exponentially decaying error from the truncation of Fourier expansion of order M of g , see [8].*

Remark 3.3 *The result obtained in Theorem 3.1 can not be improved in the exponential growth 2^M for a priori convergence analysis of general parametric functions. In fact, it can be proved that $\|g - \mathcal{I}_M[g]\|_{L^\infty(V_x)} \geq (1-\varepsilon)2^M d_M(L^\infty(V_x))$ for arbitrary small $\varepsilon > 0$ under certain assumptions [3].*

4 Applications

In this section, we study the accuracy and efficiency of the weighted empirical interpolation method (wEIM) compared to the conventional empirical interpolation method (EIM) as well as the stochastic collocation method (SCM) for one dimensional problem and sparse grid stochastic collocation method (SG-SCM) [15] for multidimensional problem. Given a function g , we denote by g_M its approximation using M “elements” (either basis functions for wEIM and EIM, or interpolation nodes for SCM and SG-SCM) and we define the error in the following two norms

$$\|g - g_M\|_{L^\infty(D; C^0(\Gamma))} \quad \text{and} \quad \|\mathbb{E}[g] - \mathbb{E}[g_M]\|_{L^\infty(D)}, \tag{4.1}$$

where the expectation $\mathbb{E}[g]$ is computed by Gauss quadrature formula specified when in need.

4.1 Parametric function in one dimension - geometric Brownian motion

We consider a geometric Brownian motion S_t satisfying a stochastic ordinary differential equation $dS_t = kS_t dt + \sigma S_t dB_t$ (This is, e.g., the most widely used model of stock price S_t at time t with drift k , volatility σ and standard Brown motion B_t [16]). The solution is given by $S_t = \exp(\sigma B_t + (k - \sigma^2/2)t)$. For simplicity, we set $S_0 = 1$, $\sigma = 1$ and $k = 1/2$ so that S_t can be written as $S_t = \exp(\sqrt{t}B_1)$, where B_1 is a standard Gauss random variable $B_1 \sim \mathcal{N}(0, 1)$. By denoting $x \equiv t$, $\mu \equiv B_1 \in \mathbb{R}^K$, $K = 1$ and $g = S_t$, we seek the following affine expansion by wEIM given in Algorithm 1

$$g(x, \mu) = \exp(\sqrt{x}\mu) \approx g_M(x, \mu) = \sum_{j=1}^M \Theta_j(\mu) q_j(x) \text{ where } \mu \sim \mathcal{N}(0, 1). \quad (4.2)$$

Moreover, we are interested in the expectation of g at time x , which can be approximated by Gauss-Hermite quadrature with abscissas and weights (μ_n, w_n) , $1 \leq n \leq N$

$$\mathbb{E}_\mu[g](x) \approx \sum_{j=1}^M \left(\int_{-\infty}^{\infty} \Theta_j(\mu) \rho(\mu) d\mu \right) q_j(x) \approx \sum_{j=1}^M \left(\sum_{n=1}^N \Theta_j(\mu_n) w_n \right) q_j(x), \quad (4.3)$$

where ρ is standard normal density function. The advantage of (4.3) is that we do not need to compute the function g for μ_n , $1 \leq n \leq N$ at every x but only at the empirical interpolation nodes x^m , $1 \leq m \leq M$, which is attributed to solving a small linear system (2.3) for $\Theta_j(\mu_n)$, $1 \leq j \leq M$, $1 \leq n \leq N$. When the evaluation of the function itself at (x, μ) is expensive and we have a large number of points x , the wEIM can be employed for efficient computation of the statistics. We set the tolerance as $\varepsilon_{tol} = 1 \times 10^{-12}$, take 1000 equidistant points in the vertex set V_x and 1000 normal distributed samples in the sample set Ξ_μ , we also take an independent 1000 normal distributed samples to test different interpolation methods. The weight in Algorithm 1 is taken as the normalized Gauss density function $w(\mu) = \rho(\mu)/\rho(0)$. As for the evaluation of the expectation of $\mathbb{E}[g_M]$, we use 12 quadrature abscissas in (4.3), which is sufficiently accurate for this example. We examine the convergence of “EIM bound” and “wEIM bound” ($r_M(x^M)$), error by “EIM test” and “wEIM test” (error computed from test samples) and test error by stochastic collocation method “SCM test”.




Figure 4.1: Comparison of convergence property of EIM, wEIM and SCM in different norms. Left: decreasing of the error $\|g - g_M\|_{L^\infty(D; C^0(\Gamma))}$; Right: decreasing of the error $\|\mathbb{E}[g] - \mathbb{E}[g_M]\|_{L^\infty(D)}$.

The convergence property of different methods is displayed in Figure 4.1, from which we can see that all the methods achieve exponential convergence rate and wEIM converges faster than both SCM and EIM in $L^\infty(D; C^0(\Gamma))$ norm. However, as for the expectation in $L^\infty(D)$ norm, SCM is the best and

wEIM is evidently better than EIM which does not take the weight into consideration. The reason for these results is that wEIM and EIM select the samples by $L_w^\infty(\Xi_\mu)$ and $L^\infty(\Xi_\mu)$ optimization, leading to small error in $L^\infty(D; C^0(\Gamma))$ norm and relatively large error for the evaluation of expectation.

4.2 Parametric function in multidimension - Karhunen-Loève expansion

For the case of multidimensional parameters, we consider the function g truncated from Karhunen-Loève expansion of a Gaussian random field with correlation length L and eigenvalues $\lambda_n, 1 \leq n \leq N_t$, written as [15]

$$g(x, \mu) - g_0(x) = C \exp \left(\left(\frac{\sqrt{\pi}L}{2} \right)^{\frac{1}{2}} \mu_1(\omega) + \sum_{n=1}^{N_t} \sqrt{\lambda_n} (\sin(n\pi x)\mu_{2n}(\omega) + \cos(n\pi x)\mu_{2n+1}(\omega)) \right) \quad (4.4)$$

where $\mu_i \sim \mathcal{N}(0, 1), 1 \leq i \leq 2N_t + 1$ are standard Gauss random variables defined in sample space $\Omega \ni \omega$. This function is widely used, e.g. in modelling the random property of porous medium in material science, geophysics, etc. To compare the convergence properties of different methods, we take $g_0 = 0, C = \exp(5), N_t = 2, L = 1/8$ and $\lambda_1 = 0.213, \lambda_2 = 0.190; x \in [0, 1]$ is discretized by 1000 equidistant vertices. We set tolerance $\varepsilon_{tol} = 1 \times 10^{-12}$, and use 1000 five dimensional independent normal distributed samples and another 1000 test samples. For the computation of $\mathbb{E}[g]$, we apply SG-SCM based on Gauss-Hermite quadrature with the deepest interpolation level 4 in each dimension.




Figure 4.2: Comparison of convergence property of EIM, wEIM and SG-SCM in different norms. Left: decreasing of the error $\|g - g_M\|_{L^\infty(D; C^0(\Gamma))}$; Right: decreasing of the error $\|\mathbb{E}[g] - \mathbb{E}[g_M]\|_{L^\infty(D)}$.

Figure 4.2 depicts the convergence rate of different methods, from which we can observe that in multidimensional problems wEIM and EIM perform much better than SG-SCM in both $\|g - g_M\|_{L^\infty(D; C^0(\Gamma))}$ error and $\|\mathbb{E}[g] - \mathbb{E}[g_M]\|_{L^\infty(D)}$ error. Both wEIM and EIM achieve fast exponential convergence rate and considerably alleviate the “curse-of-dimensionality” suffered by SG-SCM. wEIM uses only 29 samples while EIM needs 80 samples and thus 80 expansion terms, which is far less efficient than the weighted type in practical applications, e.g., in approximating the non-affine terms of reduced basis method.

4.3 Parametric equation - application in non-affine reduced basis method

As mentioned before, EIM was originally developed to deal with non-affine terms in reduced basis discretization of partial differential equations in [1]. The efficiency of the reduced basis method depends critically on the number of affine terms for both offline construction and online evaluation [10, 6, 11, 14]. Therefore, wEIM is more suitable for reduced basis approximation of non-affine parametric equation with weighted parameters.

We consider the following elliptic equation with random coefficient and homogeneous Dirichlet boundary condition: find $u : \bar{D} \times \Omega \rightarrow \mathbb{R}$ such that

$$-\nabla(g(x,\omega)\nabla u(x,\omega)) = f(x) \quad (x,\omega) \in D \times \Omega, \quad (4.5)$$

where the random coefficient $g(x,\omega)$ is a Gauss random field represented by a truncated Karhunen-Loève expansion as in (4.4). We set $D = (0,1)^2$, $f = 1$, $g_0 = 0.1$, $C = \exp(5)$, $L = 1/16$, $N_t = 5$, $\lambda_1 = 0.110$, $\lambda_2 = 0.107$, $\lambda_3 = 0.101$, $\lambda_4 = 0.095$, $\lambda_5 = 0.087$, and identify the eigenfunctions in (4.4) as $\sin(n\pi x_1)$ and $\cos(n\pi x_2)$ with $x_1, x_2 \in [0,1]$. The tolerance for weighted empirical interpolation method is taken as $\varepsilon_{tol} = 1 \times 10^{-12}$. Note that the problem has 11 independent and normal distributed random variables $\mu_K \sim \mathcal{N}(0,1)$, $1 \leq K \leq 11$ and all the random variables have relatively equivalent importance due to very close eigenvalues. Therefore, we employ isotropic sparse grid stochastic collocation method based on Gauss-Hermite quadrature [15] for the computation of statistics.

We first run wEIM and EIM with finite element vertices $|V_x| = 185$ and normal distributed samples $|\Xi_\mu| = 10000$ to build an affine expansion 1.1 for the coefficient g of problem (4.5). Another independent 1000 normal distributed samples are used to test the accuracy of the two expansions. The results are shown on the left of Figure 4.3, from which we can observe that wEIM is much more efficient with only 31 affine terms than EIM requiring 94 terms to achieve the same approximation accuracy in $L^\infty(D; C^0(\Gamma))$ norm.




Figure 4.3: Convergence property of wEIM in reduced basis approximation. Left: decreasing of the error $\|g - g_M\|_{L^\infty(D; C^0(\Gamma))}$ for EIM and wEIM; Right: decreasing of the error $\|s - s_{N,M}\|_{L^\infty(\Gamma)}$.

We use the affine expansion constructed by wEIM to build a weighted reduced basis approximation (introduced in [7] for stochastic problems) with finite element discretization in physical domain D to the stochastic elliptic problem (4.5). The quantity of interest is the integral of the solution over the physical domain D , $s = \int_D u dx$, which is computed from the finite element solution. We denote $s_{N,M}$ the approximation of s based on using N reduced bases and M affine terms. The convergence of $\|s - s_{N,M}\|_{L^\infty(\Gamma)}$ is displayed on the right of Figure 4.3, which demonstrates that wEIM is efficient in the application to reduced basis method resulting in only a few elements in the reduced basis space. Moreover, we can see that the accuracy of wEIM, represented by different number of affine terms $M = 1, 11, 21, 31$, is clearly influential to the accuracy of the reduced basis approximation.

Finally, we compare the proposed approach, a combination of weighted empirical interpolation with weighted reduced basis approximation (wEIM-RBM), to one of the most efficient stochastic computational methods - SG-SCM [15] for their accuracy and efficiency. The result of this comparison, for the $\|s - s_{N,M}\|_{L^\infty(\Gamma)}$ norm, is depicted on the left of Figure 4.4, from which the ‘‘curse-of-dimensionality’’ of SG-SCM can be obviously observed. In contrast, wEIM-RBM effectively alleviates this computational burden, using merely 15 bases to accurately approximate the stochastic solution depending on 11 independent normal distributed random variables.

As for the approximation of expectation $\mathbb{E}[s]$, we only need to compute the quantity $s_{N,M}$ with $N = 15, M = 31$ by online evaluation of reduced basis method at the sparse Gauss quadrature




Figure 4.4: Comparison of the convergence property between methods wEIM-RBM and SG-SCM. Left: decreasing of the error $\|s - s_{N,M}\|_{L^\infty(\Gamma)}$; Right: decreasing of the error $|\mathbb{E}[s] - \mathbb{E}[s_{N,M}]|$.

abscissas and then $\mathbb{E}[s_{N,M}]$ by sparse grid Gauss quadrature formula.

The comparison of wEIM-RBM with SG-SCM on the right of Figure 4.4 shows that in order to achieve the same accuracy, it takes 7 bases by reduced basis approximation while 2575 collocation nodes for stochastic collocation approximation. It is worth to mention that the online evaluation of the reduced basis method is independent of the degree of freedom ($|V_x|$) of the deterministic system. Therefore, when solving the underlying deterministic system is computational demanding (with large $|V_x|$) and the dimension of the stochastic space becomes high (with more random variables), wEIM-RBM is much more efficient than SG-SCM for non-affine stochastic problems, see [6] for detailed comparison of computational cost.

5 Concluding remarks

In order to approximate parametric functions with weighted parameters, e.g. random variables with various probability distributions, we extended the empirical interpolation method by taking the weight into account for the construction of interpolation formula. A priori convergence analysis of the weighted empirical interpolation method was provided. We obtained a direct comparison of the interpolation error to the Kolmogorov N-width, which improved the result obtained recently in [13].

By the applications in approximating geometric Brownian motion in one dimension and exponential Karhunen-Loève expansion in multidimension, we demonstrated numerically the exponential convergence rate of the weighted empirical interpolation method and its advantage in accuracy and efficiency over the empirical interpolation method as well as over the sparse grid stochastic collocation method. We also applied the proposed method to the weighted reduced basis approximation [7] for non-affine stochastic elliptic equation and illustrated its efficiency and especially its effectiveness in alleviating the “curse-of-dimensionality” in comparison with the sparse grid stochastic collocation method.

The weighted empirical interpolation method can be straightforwardly applied to nonlinear stochastic partial differential equations with reduced basis approximation and can also be employed effectively in various fields embracing weighted parameters or random variables, e.g. image science, geophysics, mathematical finance, material science, bioengineering and uncertainty quantification at large.

Acknowledgement: We acknowledge the use of the Matlab packages *MLife* previously developed by Prof. Fausto Saleri from MOX, Politecnico di Milano. We thank the anonymous referees for help insights. This work is partially supported by Swiss National Science Foundation under grant N.200021.141034. G. Rozza acknowledges the support provided by the program NOFYSAS (New opportunities for young scientists) at SISSA, International School for Advanced Studies, Trieste.

References

- [1] M. Barrault, Y. Maday, N.C. Nguyen, and A.T. Patera. An empirical interpolation method: application to efficient reduced-basis discretization of partial differential equations. *Comptes Rendus Mathématique, Analyse Numérique*, 339(9):667–672, 2004.
- [2] R. Becker, and R. Rannacher. An optimal control approach to a posteriori error estimation in finite element methods. *Acta Numerica*, 10(1):1–102, 2001.
- [3] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence rates for greedy algorithms in reduced basis methods. *SIAM Journal of Mathematical Analysis*, 43(3):1457–1472, 2011.
- [4] S. Chaturantabut and D.C. Sorensen. Nonlinear model reduction via discrete empirical interpolation. *SIAM Journal on Scientific Computing*, 32(5):2737–2764, 2010.
- [5] P. Chen, and A. Quarteroni. Accurate and efficient evaluation of failure probability for partial differential equations with random input data. *Computer Methods in Applied Mechanics and Engineering*, 267(0):233–260, 2013.
- [6] P. Chen, A. Quarteroni, and G. Rozza. Comparison of reduced basis method and collocation method for stochastic elliptic problems. *Journal of Scientific Computing*, in press, DOI 10.1007/s/10915-013-9764-2, 2013.
- [7] P. Chen, A. Quarteroni, and G. Rozza. A weighted reduced basis method for elliptic partial differential equations with random input data. *accepted in SIAM Journal on Numerical Analysis*, 2013.
- [8] R.A. DeVore and G.G. Lorentz. *Constructive Approximation*. Springer, 1993.
- [9] M.B. Giles, and E. Süli. Adjoint methods for PDEs: a posteriori error analysis and postprocessing by duality. *Acta Numerica*, 11(1):145–236, 2002.
- [10] M.A. Grepl, Y. Maday, N.C. Nguyen, and A.T. Patera. Efficient reduced-basis treatment of nonaffine and nonlinear partial differential equations. *ESAIM: Mathematical Modelling and Numerical Analysis*, 41(03):575–605, 2007.
- [11] T. Lassila, A. Manzoni, and G. Rozza. On the approximation of stability factors for general parametrized partial differential equations with a two-level affine decomposition. *ESAIM: Mathematical Modelling and Numerical Analysis*, 46:1555–1576, 2012.
- [12] T. Lassila and G. Rozza. Parametric free-form shape design with PDE models and reduced basis method. *Computer Methods in Applied Mechanics and Engineering*, 199(23):1583–1592, 2010.
- [13] Y. Maday, N.C. Nguyen, A.T. Patera, and G.S.H. Pau. A general, multipurpose interpolation procedure: the magic points. *Communications on Pure and Applied Analysis*, 8(1):383–404, 2009.
- [14] A. Manzoni, A. Quarteroni, and G. Rozza. Model reduction techniques for fast blood flow simulation in parametrized geometries. *International Journal for Numerical Methods in Biomedical Engineering*, 28(6-7):604–625, 2012.
- [15] F. Nobile, R. Tempone, and C.G. Webster. A sparse grid stochastic collocation method for partial differential equations with random input data. *SIAM Journal on Numerical Analysis*, 46(5):2309–2345, 2008.
- [16] B. Øksendal. *Stochastic Differential Equations: An Introduction with Applications*. Springer, 2010.
- [17] A. Pinkus. *N-widths in Approximation Theory*. Springer, 1985.

- [18] A. Quarteroni, G. Rozza, and A. Manzoni. Certified reduced basis approximation for parametrized partial differential equations and applications. *Journal of Mathematics in Industry*, 1(1):1–49, 2011.
- [19] A. Quarteroni, R. Sacco, and F. Saleri. *Numerical Mathematics*. Springer, 2007.
- [20] G. Rozza. Reduced basis methods for Stokes equations in domains with non-affine parameter dependence. *Computing and Visualization in Science*, 12(1):23–35, 2009.
- [21] G. Rozza, D.B.P. Huynh, and A.T. Patera. Reduced basis approximation and a posteriori error estimation for affinely parametrized elliptic coercive partial differential equations. *Archives of Computational Methods in Engineering*, 15(3):229–275, 2008.
- [22] K. Urban and B. Wieland. Affine decompositions of parametric stochastic processes for application within reduced basis methods. In *Proceedings MATHMOD, 7th Vienna International Conference on Mathematical Modelling (accepted)*, 2012.

Recent publications :

MATHEMATICS INSTITUTE OF COMPUTATIONAL SCIENCE AND ENGINEERING
Section of Mathematics
Ecole Polytechnique Fédérale
CH-1015 Lausanne

- 40.2012** F. NEGRI, G. ROZZA, A. MANZONI, A. QUARTERONI:
Reduced basis method for parametrized elliptic optimal control problems
- 41.2012** A. ABDULLE, A. BARTH, X. SCHWAB:
Multilevel Monte Carlo methods for stochastic elliptic multiscale PDES
- 42.2012** P. CHENG, A. QUARTERONI, G. ROZZA:
Uncertainty quantification of human arterial network
- 43.2012** M. ASTORINO, F. CHOULY, A. QUARTERONI:
Multiscale coupling of finite element and lattice Boltzmann methods for time dependent problems
- 44.2012** D. KRESSNER, X. LIU:
Structured canonical forms for products of (skew-)symmetric matrices and the matrix equation $XAX = B$
- 45.2012** B. BECKERMANN, D. KRESSNER, C. TOBLER:
An error analysis of Galerkin projection methods for linear systems with tensor product structure
- 46.2012** J. BECK, F. NOBILE, L. TAMELLINI, R. TEMPONE:
A quasi-optimal sparse grids procedure for groundwater flows
- 47.2012** A. ABDULLE, Y. BAI:
Fully discrete analysis of the heterogeneous multiscale method for elliptic problems with multiple scales
- 48.2012** G. MIGLORATI, F. NOBILE, E. VON SCHWERIN, R. TEMPONE:
Approximation of quantities of interest in stochastic PDES by the random discrete L^2 projection on polynomial spaces
- 01.2013** A. ABDULLE, A. BLUMENTHAL:
Stabilized multilevel Monte Carlo method for stiff stochastic differential equations
- 02.2013** D. N. ARNOLD, D. BOFFI, F. BONIZZONI:
Tensor product finite element differential forms and their approximation properties
- 03.2013** N. GUGLIELMI, D. KRESSNER, C. LUBICH:
Low-rank differential equations for Hamiltonian matrix nearness problems
- 04.2013** P. CHEN, A. QUARTERONI, G. ROZZA:
A weighted reduced basis method for elliptic partial differential equations with random input data
- 05.2013** P. CHEN, A. QUARTERONI, G. ROZZA:
A weighted empirical interpolation method: a priori convergence analysis and applications